

Delta Dental of Virginia Clinical Policy # 703

Subject Frenulectomy (Frenectomy or Frenotomy) and Frenuloplasty	Originating Department Clinical Professional Services
	Signature Authority Dental Director

Type: New Replacement Revision Clarification

Date: 11/09/2009 **Revision Date:** 11/15/2011

Preamble:	<p>The Clinical Policy Bulletin is an expression of Delta Dental of Virginia's (DDVA) determination regarding whether certain services or supplies are medically or dentally necessary. DDVA bases its conclusions on a review of currently available clinical literature. This includes, but is not limited to, clinical outcome studies published in the peer-reviewed medical and dental literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians and dentists practicing in relevant clinical areas, and other relevant factors. DDVA reserves the right to revise these policies as new clinical information is available and we welcome submission of further relevant information.</p> <p>A group may define covered dental services under their dental plan, as well as those services that may be subject to dollar caps or other limits. The plan documents outline covered benefits, exclusions and limitations. DDVA advises dentists and enrollees to consult the plan documents to determine if there are exclusions or other benefit limitations applicable to the service request. The conclusion that a particular service is medically or dentally necessary does not constitute an indication or warranty that the service requested is a covered benefit payable by DDVA. Some plans exclude coverage for services that DDVA considers either medically or dentally necessary. When there is a discrepancy between DDVA's clinical policy and the group's plan documents, DDVA is to defer to the group's plan documents as to whether the dental service is a covered benefit. In addition, if state or federal regulations mandate coverage then DDVA will adhere to the applicable regulatory requirement.</p>
History:	<p>Dentally, a frenum, or frenulum, is a mucous membrane covered connective tissue attachment that limits the movement of the lips, cheeks or tongue(1,2,13). A frenectomy (frenulectomy/frenuloplasty) is the surgical procedure utilized to correct an adverse anatomic position of a large or aberrant frenum by removal or realignment.</p> <p>Histologically, the frenum has been shown to contain dense collagen, loose connective tissue and elastic tissue fibers, but no muscle fibers. Muscle fibers may be seen in contiguous vestibular tissue but not in the frenum proper(3). A frenum may be "aberrant" in that it may develop in a disadvantageous anatomical position that adversely affects the health of other oral structures or interferes with the placement and function of tooth replacement devices. An aberrant frenum, depending on the anatomical position, is capable of retracting the gingival margin, perpetuating a diastema, affecting esthetics, limiting functional movement of the tongue, and compromising the fit of a dental prosthesis(4,5,6,7,8,9,10).</p> <p>The effect of the maxillary labial frenum, which attaches to the center of the upper lip and between the upper central incisors, on the etiology of midline diastema or</p>

	<p>space is unclear(7,11,12). A labial frenectomy removes the labial frenulum, and orthodontic patients often have this procedure done to assist in closure of a diastema or persistent space between the upper central incisors. Research has shown the probability of long-term spontaneous closure of a small diastema as the same regardless of frenectomy surgery (4) if the frenum is not very fibrous. However, large fibrous frena that obstruct diastema closure or exert a traumatic influence on the gingiva should be eliminated(4,5). There is evidence to support removal of a frenum following post-orthodontic closure of a maxillary midline diastema, as well as for instances of restrictive lip or tongue movement, and when indicated prior to placement of a prosthesis(4).</p> <p>The fit and retention of a dental prosthesis, such as a complete or removable partial denture, can be adversely affected by the presence of a frenum. Movement of a patient's lips may activate the frenum, which then, disturbingly to the patient, displaces the denture from the underlying supporting ridge. Removal of the offending frenum by frenectomy may alleviate this problem.</p> <p>The removal of a lingual frenum under the tongue can also be accomplished with either frenectomy or frenuloplasty. This surgery is indicated for patients diagnosed as being "tongue-tied." This minor surgical procedure allows increased mobility of the tongue, and the previously restricted tongue will be able to normally protrude or extend out of the mouth.</p>
Policy:	<p>DDVA Guidelines:</p> <ol style="list-style-type: none"> 1. Frenulectomy and frenuloplasty are considered incidental procedures when performed with any other surgical procedure in the same surgical area by the same dentist/dental office on the same date of service. 2. Frenectomy is appropriate for the treatment of ankyloglossia (tongue-tied). 3. Frenectomy is appropriate when the frenum develops in a disadvantageous anatomical position that adversely affects the periodontal health of other oral structures, or interferes with the placement and function of tooth replacement devices. 4. When a frenulum is attached to the alveolar ridge where it unseats a full denture prosthesis, frenectomy is appropriate to improve the stability of the denture.
Codes(14):	<p>D7960 – Frenulectomy (frenectomy or frenotomy)</p> <p>D7963 – Frenuloplasty</p>
References:	<ol style="list-style-type: none"> 1. Jablonski, Stanley. Illustrated Dictionary of Dentistry. 1982. W B Saunders Co. 2. Dorland W. Dorland's Illustrated Medical Dictionary. 30th Ed. 2004. W B Saunders Co. 3. Harris SW, Levin MP and Tsaknis PJ. Histologic features of the superior labial frenum. J Perio 1976;47:25-28. 4. Miller PD. The frenectomy combined with a laterally positioned pedicle graft. Functional and esthetic considerations. J Perio;56:102-106. 5. American Academy of Periodontology. Proceedings of the 1996 World

	<p>Workshop in Periodontics. Consensus on mucogingival therapy. <i>Annals of Periodontology</i> 1996;1;704</p> <ol style="list-style-type: none">6. Peacock ME. Frenotomy and keratinized tissue augmentation. <i>Gen Dent</i> 1998;46:194-196.7. Lindhe J, Karring T and Lang NP. <i>Clinical Periodontology and Implant Dentistry</i>. 3rd Ed. 2000. Munksgaard, Copenhagen.8. Bagga S, Bhat KM, et al. Esthetic management of the upper labial frenum: a novel frenectomy technique. <i>Quintessence Int</i> 2006;37:819-823.9. American Academy of Pediatric Dentistry. Council on Clinical Affairs. Guideline on Pediatric Oral Surgery 2005;205-211.10. McCracken's Removable Partial Prosthodontics. Carr AB, McGivney GP and Brown PT. 11th Ed. 2005. C V Mosby Co. St Louis.11. The diastema, the frenum, the frenectomy: a clinical study. Edwards JG. <i>Amer J Orthod</i> 1977;71:489-508.12. The maxillary inter-incisal diastema and its relationship to the superior labial frenum and the inter-maxillary suture. Popovich F, Thompson GW and Main PA. <i>Angle Orthod</i> 1977;47:265-271.13. American Dental Association. <i>Current Dental Terminology. CDT 2011-2012</i>; 213 (©ADA 2010).14. American Dental Association. <i>Current Dental Terminology. CDT 2011-2012</i>; 67 (©ADA 2010).
--	--